

Annual Review of Biomedical Engineering
Mechanical Control of Cell
Differentiation: Insights
from the Early Embryo

Celeste M. Nelson

Departments of Chemical & Biological Engineering and Molecular Biology, Princeton University, Princeton, New Jersey USA; email: celesten@princeton.edu

ANNUAL
REVIEWS **CONNECT**

www.annualreviews.org

- Download figures
- Navigate cited references
- Keyword search
- Explore related articles
- Share via email or social media

Annu. Rev. Biomed. Eng. 2022. 24:307–22

First published as a Review in Advance on
April 6, 2022

The *Annual Review of Biomedical Engineering* is
online at bioeng.annualreviews.org

<https://doi.org/10.1146/annurev-bioeng-060418-052527>

Copyright © 2022 by Annual Reviews. This work is licensed under a Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See credit lines of images or other third-party material in this article for license information

Keywords

morphodynamics, mechanical stress

Abstract

Differentiation is the process by which a cell activates the expression of tissue-specific genes, downregulates the expression of potency markers, and acquires the phenotypic characteristics of its mature fate. The signals that regulate differentiation include biochemical and mechanical factors within the surrounding microenvironment. We describe recent breakthroughs in our understanding of the mechanical control mechanisms that regulate differentiation, with a specific emphasis on the differentiation events that build the early mouse embryo. Engineering approaches that reproducibly mimic the mechanical regulation of differentiation will permit new insights into early development and applications in regenerative medicine.

Contents

INTRODUCTION	308
FROM OOCYTE TO ORGANISM—THE ROLE OF MECHANICS IN THE SYMPHONY OF DIFFERENTIATION EVENTS THAT BUILD THE EARLY MAMMALIAN EMBRYO	310
Cortical Tension and Pressure: The Mechanical Forces that Sculpt the Blastula . . .	310
The Most Important Stage of Your Life: The Mechanics of Gastrulation	314
LESSONS FROM THE EARLY EMBRYO	316

INTRODUCTION

From a single, fertilized egg emerges the approximately 200 or so phenotypically distinct cell types that make up a human or a mouse. These cells are generated through the process of differentiation, a cascade of events that endow cells with identity and function and permit them to construct the tissues and organs necessary for multicellular life (1, 2). Differentiation continues apace in mature tissues, wherein resident stem or progenitor cells divide and differentiate to retain or restore population levels of tissue-specific cell types during homeostasis or after injury (3). The importance of differentiation can also be appreciated by considering a variety of different diseases, including fibrosis and cancer, in which cells within tissues dedifferentiate into a more stem-like state or transdifferentiate to take on new deleterious phenotypes (4).

Our understanding of the molecular signaling pathways and extracellular stimuli that regulate differentiation has deepened over the last decade. These breakthroughs were driven in part by advances in the isolation and culture of embryonic stem (ES) cells (5) and induced pluripotent stem (iPS) cells (6); these advances revealed the transcription factors and other proteins that are essential for maintaining a cell in an undifferentiated state (7, 8), for promoting its self-renewal (9), and for inducing its differentiation down a specific lineage (10). This rich understanding of fundamental mechanisms has permitted the use of stem cells to repair damaged organs in regenerative medicine applications (5). Parallel studies using stem cells isolated from adult tissues, including mesenchymal stem cells (MSCs), showed that the mechanical properties (see the sidebar titled A Brief Primer on Mechanics in Biological Systems) of the microenvironment cooperate with—and can even override—information from biochemical signals to induce differentiation (11–13). For example, in a now-classic study, MSCs that were cultured on hydrogel-based substrata engineered to mimic the stiffness of the brain differentiated down a neurogenic lineage, whereas those cultured on substrata engineered to mimic the stiffness of bone differentiated down an osteogenic route (14). This switch in cell fate depends in part on the physical properties of the nucleus itself; these properties are regulated by scaffolding proteins such as lamin-A, the expression of which is tuned by substratum stiffness (15). Similarly, experiments using Transwell filters with different pore sizes revealed that migration through confined spaces enhances the differentiation of MSCs down the osteogenic lineage (16). The concept of mechanical regulation of stem cell differentiation was thus pioneered through the clever use of engineered culture microenvironments.

Studies focused on the differentiation of individual cells in culture likewise permitted the creation of three-dimensional (3D) *in vitro* mimetics of tissues and organs known as organoids (17–19). The first stem cell–derived organoids reproduced the 3D organizations of the crypt–villus axis of the intestine (20) and the cortex of the brain (21). In the intervening years, stem cells have been differentiated to form 3D organoids that mimic the architectures of a variety of organs, including different regions of the brain, gastrointestinal tract, liver, pancreas, lung, and kidney (reviewed in

ES: embryonic stem

iPS: induced pluripotent stem

A BRIEF PRIMER ON MECHANICS IN BIOLOGICAL SYSTEMS

Cells and tissues, including those of the developing embryo, must obey Newton's laws of motion. To understand the role of mechanical forces in differentiation, we need to consider the active inputs (the stresses, strains, and flows that impinge upon a cell), the passive inputs (the material properties of cells and their surrounding microenvironment), and the mechanical and biochemical responses of the cell itself.

- **Active inputs:** A force is any input that causes an object, such as a cell or a tissue, to change its shape (deformation) or velocity (speed and direction). In biological systems, forces result from fluid flows, fluid pressures, and tension or compression induced by other cells or tissues. Cells appear capable of sensing both stress (force per unit area, akin to pressure) and strain (deformation per unit length, akin to stretch). Although force and stress are related to each other and often used interchangeably, these parameters are not identical. The application of a constant force can lead to levels of stress that change over space and time as the object (the cell or tissue) deforms.
- **Material properties:** The mechanical stiffness of any object, including a cell as well as its extracellular microenvironment, affects how that object will respond to the application of force. The stiffer the object, the more force is required to deform it. Compliance and elasticity are commonly used terms to describe stiffness; the softer the object, the more compliant or elastic it is. The Young's (or elastic) modulus refers to the stiffness of a material, independent of its geometry. The higher the modulus, the stiffer the material.
- **Cellular responses to mechanical forces:** Cells use their cytoskeletons and adhesive contacts to probe the mechanical properties of their surrounding microenvironment. Cytoskeletal tension refers to the pulling forces exerted by the actin, microtubule, and/or intermediate filament networks within a cell. The higher the cytoskeletal tension, the more the cell pulls on its neighbors or extracellular matrix. Cortical tension refers to the contractility of the actomyosin cytoskeleton located at the cell membrane. The higher the cortical tension, the greater the tendency for the cell to round up. Each network responds to changes in the mechanical properties or forces imparted on a cell through processes known as mechanosensing and mechanotransduction.

17 and 22). Stem cell–derived organoids have been used to model human diseases, both for studying basic mechanisms and for screening of potential therapeutics (23, 24). These are exciting applications of a technology borne out of insights into the fundamental processes of differentiation. Furthermore, organoids have served as minimal culture models that mimic aspects of embryonic development and have allowed scientists and engineers to uncover the step-wise ordering of extracellular signals and gene-expression patterns that trigger a pluripotent or multipotent cell to differentiate down distinct lineages outside of the embryo (25, 26).

As with isolated stem cells in two-dimensional (2D) culture, 3D organoid models have also started to reveal the relative roles of mechanical forces in the regulation of differentiation. For example, synthetic matrices that mimic the stiffness of the liver promote the proliferation of liver progenitor cells into organoids (27), whereas matrices that are softer than the liver prevent organoid formation. In contrast, similar experiments with intestinal progenitor cells revealed that the size of the stem cell compartment within intestinal organoids decreases as stiffness increases (28). How stem cells interpret the mechanical properties of their local microenvironment as they form into organoids is therefore likely dependent on their tissue type.

Insights from the above-described culture models, both 2D and 3D, have ignited interest in defining the relative roles of mechanical forces and mechanical signaling in the regulation of differentiation in embryonic and fetal tissues *in vivo*. It is possible that many of the same regulatory

Blastocyst: an early stage of embryonic development in mammals that consists of a hollow ball of cells surrounding an inner cell mass

Inner cell mass (ICM): the cells of the blastocyst that are fated to become the embryo itself along with the primitive endoderm

Trophectoderm (TE): the cells at the outer layer of the blastocyst that are fated to become part of the placenta

Morula: a solid ball of cells generated from early cleavage divisions of a zygote

mechanisms that control differentiation in culture will be found to operate in embryonic and fetal tissues; however, it is just as (if not more) likely that the clever reductionist approaches described above overlook key complexities of the embryonic microenvironment and that an entirely different set of regulatory mechanisms will be discovered from studies *in vivo*. This review serves as a primer for early embryonic development of the mouse and highlights the mechanical signaling pathways and mechanical forces that have so far been found to instruct each differentiation event that specifies cell fate in the early embryo. We begin with a discussion of the preimplantation embryo, at a stage when each of the eight cells that compose the developing mouse is essentially identical with respect to its differentiation potential. We then describe the mechanical signaling involved in symmetry breaking, segregation of the embryonic and extraembryonic tissues, implantation into the uterus, and differentiation of the three germ layers.

FROM OOCYTE TO ORGANISM—THE ROLE OF MECHANICS IN THE SYMPHONY OF DIFFERENTIATION EVENTS THAT BUILD THE EARLY MAMMALIAN EMBRYO

Cortical Tension and Pressure: The Mechanical Forces that Sculpt the Blastula

In the preimplantation mammalian (mouse) embryo, the first differentiation event occurs after the 8-cell stage (**Figure 1a**), when the cells of the blastocyst generate the first lineages that will become the inner cell mass (ICM), which gives rise to the embryo proper, and the outer extraembryonic trophectoderm (TE), which gives rise to the placenta. Mammalian oocytes are spherical and therefore lack any obvious geometric polarity signal that would influence later symmetry-breaking and differentiation decisions (29–31); thus, the cells of the 8-cell-stage embryo are equivalent in their geometry as well as their developmental potential. However, asymmetric division and subsequent cellular rearrangements generate a 16-cell morula that contains a core of nonpolarized cells surrounded by a layer of cells with apicobasal polarity (32). This geometric arrangement appears to require differences in myosin-mediated contraction of the apical actin cortex, such that cells with higher cortical tension become internalized and surrounded by those with lower cortical tension (33–36). It remains unclear how cortical tension is specified within cells at the 8-cell stage, but differences in the distributions of proteins at the apical surfaces of these cells are likely involved (36). Regardless of the underlying molecular details, it is notable that the first symmetry-breaking event in the mammalian embryo is generated by mechanical forces resulting from cortical tension.

This symmetry-breaking event sets up the 16-cell- and 32-cell-stage murine embryos, both of which contain a single layer of outer cells that have apicobasal polarity around a core of nonpolarized inner cells. In the first lineage-specification event, the nonpolarized inner cells become the ICM while the outer cells become the TE. Mechanical signaling also appears to be critical for driving this fate specification (**Figure 1b**). In particular, subcellular localization of Yes-associated protein (Yap) distinguishes between the two fates. Yap functions as part of the Hippo signaling pathway (**Figure 2**): When the Hippo pathway is activated, the serine/threonine kinases Lats1 and Lats2 phosphorylate Yap, causing its cytoplasmic retention (37, 38). When Hippo is inactive, Yap shuttles to the nucleus and interacts with Tead-family transcription factors to activate gene expression (39). In parallel, Yap localization and activity are also regulated by mechanical forces (40). Specifically, myosin-mediated contraction of F-actin appears to induce and sustain nuclear localization of Yap and downstream gene expression (41–43). As a consequence, mechanical forces and features of the microenvironment that are generally associated with the formation of actin stress fibers and cell spreading—including shear forces, mechanical compression, and local microenvironmental stiffness—are also associated with the nuclear localization of Yap and activation of gene expression (44, 45).




Figure 1

Mechanical forces from cortical tension and luminal fluid pressure influence differentiation in the early mouse embryo. (a) At the 8-cell stage, all cells of the embryo are equally capable of differentiating into the different cell types of the embryo proper and the extraembryonic tissues. (b) At the 16-cell stage, the embryo contains two layers of cells—an inner layer with high cortical tension and low nuclear Yap that becomes the ICM, and an outer layer with low cortical tension and high nuclear Yap that becomes the TE. (c) At the 32-cell stage, the early blastocyst develops a fluid-filled cavity that localizes to one side of the embryo and distinguishes the mural TE from the polar TE, which is adjacent to the ICM. Abbreviations: ICM, inner cell mass; TE, trophectoderm; Yap, Yes-associated protein.

In a manner consistent with the spatial patterns of cellular distention within the early mouse embryo, Yap is phosphorylated and retained in the cytoplasm in the inner cells, whereas Yap is localized to the nucleus in the outer cells (46–48). As soon as spatial differences in Yap localization are detectable, the expression of the TE-associated transcription factor *Cdx2* begins to correlate with nuclear Yap (48). *Cdx2* represses the pluripotency factors *Oct4* and *Nanog* (47, 49), thus limiting their expression (and pluripotency) to the cells of the ICM. In parallel, Yap represses the expression of the pluripotency marker *Sox2* in the outer cells, thus initiating the cascade of events that leads these cells to differentiate into TE (50). Intriguingly, the Yap cofactor *Tead4* is also responsible for a key metabolic switch in the two populations of cells at this stage of development: The ICM produces ATP largely through glycolysis (51, 52), while *Tead4*-induced signaling within the TE promotes the expression of mitochondrial transport chain components and the production of ATP through oxidative phosphorylation (53). Consistently, disrupting *Tead4* expression (54, 55) or inhibiting Rho kinase (50) causes the outer layer of cells to express ICM markers and thus prevents their differentiation into the TE.




Figure 2

Mechanical signaling through the Hippo/Yap pathway. Activation of the Hippo pathway induces phosphorylation of Lats1/2, which phosphorylate Yap and cause its retention in the cytoplasm.

The second lineage-specification event occurs during the early blastocyst (32-cell) stage (**Figure 1c**), when ICM cells begin to express markers of either epiblast or primitive endoderm fate in a stochastic manner (56–58), influenced by their relative position. Specifically, epiblast cells are biased to form adjacent to the TE and are separated from the developing lumen by the primitive endoderm (59–61). At the same time as these fates are specified, secretion of cytoplasmic vesicles causes fluid droplets to accumulate between the inner cells, which eventually coalesce or coarsen to form one large fluid-filled cavity that compresses the ICM to one side of the embryo (62–64). The pressure of this fluid was recently found to increase over the course of embryonic development (65) and to alter the relative numbers of cells in the ICM and the TE in a tension-dependent manner (66). Higher pressure leads to increased expansion of the lumen, and the forces from expansion are transmitted into maturation of tight junctions to maintain blastocyst integrity. Lower pressure appears to result in decreased luminal expansion and an increase in asymmetric divisions, favoring the generation of cells that populate the ICM. In this case, a mechanical signal (pressure) regulates fate decisions by altering the pattern of cell division within the early embryo. Although not shown explicitly, higher pressures would also be expected to increase the strain and stretch within the TE cell layer and thus enhance nuclear Yap and expression of Cdx2. As the lumen expands, the epiblast and primitive endoderm cells appear to undergo differential sorting into their final positions within the ICM (64), with those cells biased toward primitive endoderm fate moving directionally toward the ICM–lumen interface (64). Decreasing luminal pressure mechanically or biochemically decreases specification of primitive endoderm and sorting

Epiblast:

the outermost layer of cells in an embryo that will later give rise to ectoderm and mesoderm

Primitive endoderm:

a layer of cells in the early embryo that gives rise to the yolk sac

from the epiblast population (64). Whether this sorting is regulated predominantly by chemotaxis, durotaxis, or some other mechanism remains to be determined.

At the mid-blastocyst stage, the cells within the TE have already segregated into two distinct populations with completely different fates—the polar TE, which directly contacts the ICM, and the mural TE, which lacks these heterotypic contacts but surrounds the fluid-filled blastocoel. At the late-blastocyst stage (*E4.5* in the mouse), the mural TE contacts the endometrium of the uterus and initiates the process of implantation, which leads to drastic changes in the shape and potency of the epiblast. The initially oval, naive epiblast is constricted by its surrounding basement membrane as it grows, forcing it to adopt a circular shape (67, 68). Increased tension and apical constriction within the polar TE exerts force on the epiblast; together, the polar TE and epiblast transform into the postimplantation morphology known as the egg cylinder (67). These changes occur concomitantly with transition of the epiblast from naive to primed pluripotency, but it remains unclear whether the forces of remodeling are correlative or causal in this transition.

After the initiation of implantation, the highly invasive TE cells stop undergoing cytokinesis and commence endoreduplication, which generates highly polyploid primary trophoblast giant cells (TGCs) (69). Primary TGCs promote remodeling of the endometrial tissue by secreting cathepsins and matrix metalloproteinases (MMPs) (70). These cells remove adjacent endometrial epithelial cells via phagocytosis, invade into the uterine stroma, and establish connections with the maternal vasculature (71). Differentiation of TGCs appears to involve an epithelial-mesenchymal transition (EMT), in which cell–cell interactions are destabilized as cells increase their motility and invasiveness and engulf the maternal epithelial and blood cells (71, 72). EMT is then followed by reepithelialization of the postmitotic giant cells to form the vascular structures of the placenta (71, 73). Both EMT and EMT-associated failure of cytokinesis are known to be regulated by mechanical signaling in adult epithelial cells (see the sidebar titled Mechanical Regulation of Epithelial-Mesenchymal Transition and Endoreduplication). It is therefore possible that the mechanical properties of the microenvironment of the endometrium are partly responsible for initiating differentiation of TGCs. Consistent with this possibility, alterations in the mechanical properties of the endometrium were recently found to promote implantation (74).

Endoreduplication: a specialized cell cycle in which the cell replicates its genome without undergoing mitosis, thus resulting in polyploidy

TGC: trophoblast giant cell

MMP: matrix metalloproteinase

EMT: epithelial-mesenchymal transition

MECHANICAL REGULATION OF EPITHELIAL-MESENCHYMAL TRANSITION AND ENDOREDUPPLICATION

Epithelial-mesenchymal transition (EMT) is a prototypic example of transdifferentiation that is essential for development of the embryo proper and is also implicated in fibrosis and metastasis (75, 76). EMT is typically stimulated by soluble signals including transforming growth factor- β (TGF β) and matrix metalloproteinases (MMPs) and is initiated by changes in the expression of key transcription factors including members of the Snail, Twist, and ZEB families (77, 78). These transcription factors, in turn, lead to the repression of epithelial markers, including E-cadherin, and the expression of mesenchymal cytoskeletal proteins such as vimentin. These molecular changes promote the weakening of cadherin-associated intercellular adhesions and enhance single-cell migration and invasion.

As a transdifferentiation process, EMT is also regulated by the mechanical properties of the surrounding microenvironment. For example, the ability of signals such as TGF β or MMPs to induce EMT depends on the mechanical stiffness of the underlying matrix. Specifically, exposing cells in stiff microenvironments to EMT inducers leads to elevated expression of Snail and robust EMT (79–83). In contrast, soft microenvironments prevent the induction of Snail, apparently due to decreased signaling through β 1 integrin (80, 84). Strikingly, cells that continue to proliferate while undergoing EMT in culture fail to undergo cytokinesis and become multinucleated (85–87), a process that is also promoted by stiff microenvironments and inhibited by soft microenvironments (86).

The polar TE cells that are located adjacent to the ICM continue to undergo cytokinetic divisions and generate the remaining populations of cells that form the placenta (88).

In the human embryo, development and cell-fate specification are likely regulated by different mechanical events than those described above for the mouse. Compaction and cavitation occur later in the human embryo, at the 16-cell and 64-cell stages, respectively (89), and thus the mechanical forces experienced by individual cells at these stages will be different. More strikingly, Yap is localized to the nucleus in both the ICM and TE cells of the preimplantation human embryo (90), suggesting that signaling from cytoskeletal tension and/or Hippo is not responsible for this cell-fate decision in the human. These differences between the early mouse and human embryos suggest caution in generalizing results, both between species as well as between differentiation in culture and in vivo.

The Most Important Stage of Your Life: The Mechanics of Gastrulation

After the cells of the fertilized oocyte are specified as embryonic or extraembryonic, these cells then undergo a complex and highly coordinated series of rearrangements to transform from a single layer of tissue into three apposed layers of endoderm, mesoderm, and ectoderm, known as the primary germ layers. These germ layers give rise to all of the different tissues of the mature organism. As the innermost layer, the endoderm forms the epithelial lining of the gastrointestinal tract, respiratory tract, urogenital system, and endocrine glands. The middle layer, the mesoderm, differentiates into the mesenchyme, mesothelium, striated and smooth musculature, and red blood cells. As the outermost layer, the ectoderm gives rise to the epithelium of the skin and its appendages (including the mammary and sweat glands) as well as neural tissues of the brain and spinal column. The morphogenetic process that generates the primary germ layers is known as gastrulation and varies greatly between different classes of vertebrates (91, 92).

In the mouse, gastrulation begins approximately two days after implantation. Prior to gastrulation, the cup-shaped embryo is composed of a radially symmetric layer of pluripotent cells known as the epiblast (**Figure 3**), which is surrounded laterally by a layer of extraembryonic tissue known as the visceral endoderm (VE) and proximally by the trophoblast or extraembryonic ectoderm (ExE). The ExE forms the fetal portion of the placenta while the VE forms the extraembryonic membranes. The process of implantation provides the embryo with a proximal-distal axis, with the surface of the endometrium serving as the proximal end of the structure. Induction of gastrulation requires the embryo to break its radial symmetry and establish an anterior-posterior axis (which also specifies the head and tail of the organism). At the onset of gastrulation, cells at the posterior pole of the epiblast undergo morphological changes to form the primitive streak. Cells located at the primitive streak undergo an EMT, ingress, and migrate to fill the space between the epiblast and the VE as they form the definitive mesoderm.

Similar to the preimplantation embryo, symmetry breaking of the postimplantation embryo is driven in part by mechanical forces. Prior to the onset of gastrulation, the postimplantation epiblast is surrounded by and constrained within a presumably rigid basement membrane. Small, evenly distributed perforations within the basement membrane, which would weaken the constraint, correlate with growth of the epiblast (93). At the same time as the epiblast is growing, a cluster of extraembryonic cells, known as anterior VE (AVE) cells, emerges at the distal tip of the structure and migrates collectively toward one side (94) where the cells secrete inhibitors of the Wnt and Nodal pathways to inhibit posterior cell fates and thereby establish the anterior pole of the embryo (95). This collective migration is preceded by apical constriction (96) and requires activation of the Rho GTPase Rac1, which permits the AVE cells to extend actin-rich cytoskeletal projections in the direction of migration and to exchange positions with their neighbors




Figure 3

Mechanical signals from the basement membrane along with increases in tissue fluidity promote gastrulation of the mouse embryo. MMP-driven weakening of the basement membrane (indicated by perforations) permits growth of the epiblast (*left*). AVE cells secrete Nodal inhibitors, thus blocking expression of MMPs, leading to an intact basement membrane on the anterior pole of the embryo (*center*). Cells in the posterior side of the embryo undergo EMT and migrate through the weakened basement membrane, establishing the mesoderm (*right*). Abbreviations: AVE, anterior visceral endoderm; EMT, epithelial-mesenchymal transition; ExE, extraembryonic ectoderm; MMP, matrix metalloproteinase; VE, visceral endoderm.

(97)—essentially, this population of cells becomes more fluid-like (see the sidebar titled Solid-Like Versus Fluid-Like Tissues) by altering their packing (98). Particle imaging velocimetry analysis of fluorescently tagged embryos revealed that AVE cells undergo push-pull motions as they migrate collectively, and knockout screens identified the Rho GEF beta-PIX as an essential element in localizing the protrusive activity (99).

Collective migration of the AVE cells correlates with a loss of perforations within the basement membrane, such that the perforations are preferentially skewed toward the posterior side of the epiblast (93). The perforations themselves result from Nodal-dependent activation of the expression of MMP2 and MMP14, which degrade the basement membrane at the posterior side. Since the AVE cells secrete Nodal inhibitors, MMP expression is inhibited at the anterior side and the basement membrane remains intact, thus establishing a mechanical asymmetry concomitantly with the anterior-posterior axis of the embryo. Local weakening of the basement membrane at the posterior side corresponds with formation of the primitive streak—cells within this region of the epiblast undergo an EMT and breach the weakened basement membrane as they ingress to form the mesoderm (93). Generation of the germ layers thus requires changes in the mechanical properties of the microenvironment within the developing embryo.

SOLID-LIKE VERSUS FLUID-LIKE TISSUES

By definition, epithelial tissues comprise sheets of cells that form adhesive contacts with their neighbors. The shapes of the individual cells affect how they pack together to form their resident tissues and can be described using similar conceptual parameters as the ones that physicists use to describe the packing of foams (100). However, cells and tissues are not passive objects, and recent insights into their mechanical properties have been gleaned by considering them as forms of active, granular matter. One metric that is commonly used to predict the mechanical properties and behaviors of epithelial sheets is the cell-shape index, which is given as the ratio of a cell's perimeter to the square root of its area (101–103). The cell-shape index is smallest for a cell with a circular cross section and largest for an elongated cell. Therefore, for a given cross-sectional area, an epithelial cell within a stereotypical cobblestone monolayer has minimized its connections with its neighbors and its cell-shape index. These conditions are typically associated with static epithelial tissues in which the resident cells have limited motility and maintain their neighbors, a so-called jammed or frozen, solid-like state. In contrast, an epithelial cell with an elongated morphology has increased its connections with its neighbors and its cell-shape index, conditions that are associated with migratory epithelial tissues in which the resident cells undergo neighbor exchanges in an unjammed or fluid-like state. These two extremes, solid-like and fluid-like, have different mechanical properties: The jammed tissue is more rigid and mechanically stable than the unjammed tissue, which is more plastic and malleable.

The transition between jammed and unjammed tissues is similar to a glassy phase transition and has been implicated in studies of epithelial sheets in culture as well as in developing embryos (reviewed in 104). In *Drosophila*, the unjammed state is associated with an increase in fluidity that permits the movements necessary for formation of the ventral furrow during gastrulation (105), convergent-extension during morphogenesis of the germ band (106), and cellular rearrangements in the wing disc epithelium (107). In the zebrafish, the unjammed state represents a phase transition that decreases the rigidity of the tissue and promotes movement of the blastoderm over the yolk sac during epiboly (108, 109) and elongation of the body axis (110) and tail bud (111, 112). Similarly, the unjammed state corresponds to an increase in fluidity during gastrulation in *Xenopus* embryos (113). In avian embryos, an increase in tissue fluidity promotes formation of the primitive streak in quail (114), elongation of the body axis in chicks (115), and extension of the airway epithelium in the chick lung (116).

LESSONS FROM THE EARLY EMBRYO

We often anthropomorphize the differentiation of a pluripotent cell by referring to this event as a cell-fate decision, implicitly suggesting that the cell in question is presented with a variety of options and deliberately chooses its fate according to its developmental aspirations or simple expedience. In the embryo, these cell-fate decisions often accompany a symmetry-breaking event—a population of genetically identical cells becomes heterogeneous because of changes in morphology, positioning, local mechanical properties, or differential acquisition of apicobasal polarity. Although these alterations have been historically classified by corresponding differences in the expression or localization of proteins, it is important to note that each would also be expected to lead to heterogeneities in the mechanical microenvironment: A cell with apicobasal polarity experiences different mechanical forces than a cell completely surrounded by neighbors in all three dimensions. As described above, these differences in mechanical forces can themselves alter signaling, in particular through the subcellular localization of mechanosensors such as Yap, and thereby differentially direct differentiation.

The mechanical signaling networks that regulate the symphony of differentiation events in the early embryo are also important for regulating pluripotency and differentiation of stem cells in culture. ES cells are derived from the ICM and have the potential to differentiate into all cells present in the mature organism (117). Pluripotency appears to be defined specifically by the

expression of four different transcription factors, Oct4, Sox2, Klf4, and Myc (117–119), which are also expressed in the early embryo. Differentiated adult cells can be reprogrammed into iPS cells by expressing these four factors (120). Yap is highly expressed in ES cells (121), inactivated during their differentiation (122), and enhanced during the reprogramming of iPS cells, suggesting that Yap itself (and possibly mechanical signaling) is involved in the switch to and from pluripotency.

The mechanical signals that instruct cell-fate decisions in the early embryo also appear to play instructive roles during later differentiation events, both in vivo and in culture. Symmetry breaking in intestinal organoids (which are initially spherical, just like the preimplantation embryo) is accompanied by differences in the subcellular localization of Yap, which is required to enable the Paneth cell lineage by activating Notch signaling via Dll1 (123). Regulation of luminal fluid volume (and presumably fluid pressure) is essential for differentiation of enterocytes and morphogenesis of the crypt-villus axis in intestinal organoids (124, 125). Similarly, the pressure of the luminal fluid within the embryonic mouse lung regulates the rate of morphogenesis, differentiation of resident cell types, and maturation of the organ (126, 127). For most developing systems, the mechanical signaling pathways downstream of fluid pressure remain to be uncovered, but it will be interesting to define the overlap between organogenesis in vivo, organoid morphogenesis ex vivo, and mechanotransduction in the early embryo.

DISCLOSURE STATEMENT

The author is not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

Work from the author's group was supported in part by grants from the National Institutes of Health (HL110335, HL118532, HL120142, CA187692, and CA214292) and the National Science Foundation (2124582) and by a Faculty Scholars Award from the Howard Hughes Medical Institute.

LITERATURE CITED

1. Chazaud C, Yamanaka Y. 2016. Lineage specification in the mouse preimplantation embryo. *Development* 143:1063–74
2. Wu G, Scholer HR. 2016. Lineage segregation in the totipotent embryo. *Curr. Top. Dev. Biol.* 117:301–17
3. Snippert HJ, Clevers H. 2011. Tracking adult stem cells. *EMBO Rep.* 12:113–22
4. Aydin B, Mazzoni EO. 2019. Cell reprogramming: the many roads to success. *Annu. Rev. Cell Dev. Biol.* 35:433–52
5. Murry CE, Keller G. 2008. Differentiation of embryonic stem cells to clinically relevant populations: lessons from embryonic development. *Cell* 132:661–80
6. Yamanaka S. 2012. Induced pluripotent stem cells: past, present, and future. *Cell Stem Cell* 10:678–84
7. Smith AG, Heath JK, Donaldson DD, Wong GG, Moreau J, et al. 1988. Inhibition of pluripotential embryonic stem cell differentiation by purified polypeptides. *Nature* 336:688–90
8. Sato N, Meijer L, Skaltsounis L, Greengard P, Brivanlou AH. 2004. Maintenance of pluripotency in human and mouse embryonic stem cells through activation of Wnt signaling by a pharmacological GSK-3-specific inhibitor. *Nat. Med.* 10:55–63
9. He S, Nakada D, Morrison SJ. 2009. Mechanisms of stem cell self-renewal. *Annu. Rev. Cell Dev. Biol.* 25:377–406
10. Venkei ZG, Yamashita YM. 2018. Emerging mechanisms of asymmetric stem cell division. *J. Cell Biol.* 217:3785–95
11. McBeath R, Pirone DM, Nelson CM, Bhadriraju K, Chen CS. 2004. Cell shape, cytoskeletal tension, and RhoA regulate stem cell lineage commitment. *Dev. Cell* 6:483–95

12. Perestrelo T, Correia M, Ramalho-Santos J, Wirtz D. 2018. Metabolic and mechanical cues regulating pluripotent stem cell fate. *Trends Cell Biol.* 28:1014–29
13. Chaudhuri O, Mooney DJ. 2012. Stem-cell differentiation: anchoring cell-fate cues. *Nat. Mater.* 11:568–69
14. Engler AJ, Sen S, Sweeney HL, Discher DE. 2006. Matrix elasticity directs stem cell lineage specification. *Cell* 126:677–89
15. Swift J, Ivanovska IL, Buxboim A, Harada T, Dingal PC, et al. 2013. Nuclear lamin-A scales with tissue stiffness and enhances matrix-directed differentiation. *Science* 341:1240104
16. Smith LR, Irianto J, Xia Y, Pfeifer CR, Discher DE. 2019. Constricted migration modulates stem cell differentiation. *Mol. Biol. Cell* 30:1985–99
17. Rossi G, Manfrin A, Lutolf MP. 2018. Progress and potential in organoid research. *Nat. Rev. Genet.* 19:671–87
18. Schutgens F, Clevers H. 2020. Human organoids: tools for understanding biology and treating diseases. *Annu. Rev. Patol. Mech. Dis.* 15:211–34
19. Kim J, Koo BK, Knoblich JA. 2020. Human organoids: model systems for human biology and medicine. *Nat. Rev. Mol. Cell Biol.* 21:571–84
20. Sato T, Vries RG, Snippert HJ, van de Wetering M, Barker N, et al. 2009. Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. *Nature* 459:262–65
21. Eiraku M, Watanabe K, Matsuo-Takasaki M, Kawada M, Yonemura S, et al. 2008. Self-organized formation of polarized cortical tissues from ESCs and its active manipulation by extrinsic signals. *Cell Stem Cell* 3:519–32
22. Hofer M, Lutolf MP. 2021. Engineering organoids. *Nat. Rev. Mater.* 6:402–20
23. Clevers H. 2016. Modeling development and disease with organoids. *Cell* 165:1586–97
24. Li M, Izpissua Belmonte JC. 2019. Organoids—preclinical models of human disease. Reply. *N. Engl. J. Med.* 380:1982
25. Renner M, Lancaster MA, Bian S, Choi H, Ku T, et al. 2017. Self-organized developmental patterning and differentiation in cerebral organoids. *EMBO J.* 36:1316–29
26. Takebe T, Wells JM. 2019. Organoids by design. *Science* 364:956–59
27. Sorrentino G, Rezakhani S, Yildiz E, Nuciforo S, Heim MH, et al. 2020. Mechano-modulatory synthetic niches for liver organoid derivation. *Nat. Commun.* 11:3416
28. Perez-Gonzalez C, Ceada G, Greco F, Matejic M, Gomez-Gonzalez M, et al. 2021. Mechanical compartmentalization of the intestinal organoid enables crypt folding and collective cell migration. *Nat. Cell Biol.* 23:745–57
29. Hiiragi T, Solter D. 2004. First cleavage plane of the mouse egg is not predetermined but defined by the topology of the two apposing pronuclei. *Nature* 430:360–64
30. Motosugi N, Dietrich JE, Polanski Z, Solter D, Hiiragi T. 2006. Space asymmetry directs preferential sperm entry in the absence of polarity in the mouse oocyte. *PLOS Biol.* 4:e135
31. Rossant J, Tam PP. 2004. Emerging asymmetry and embryonic patterning in early mouse development. *Dev. Cell* 7:155–64
32. Johnson MH, Ziomek CA. 1983. Cell interactions influence the fate of mouse blastomeres undergoing the transition from the 16- to the 32-cell stage. *Dev. Biol.* 95:211–18
33. Anani S, Bhat S, Honma-Yamanaka N, Krawchuk D, Yamanaka Y. 2014. Initiation of Hippo signaling is linked to polarity rather than to cell position in the pre-implantation mouse embryo. *Development* 141:2813–24
34. Maitre JL, Turlier H, Illukkumbura R, Eismann B, Niwayama R, et al. 2016. Asymmetric division of contractile domains couples cell positioning and fate specification. *Nature* 536:344–48
35. Samarage CR, White MD, Alvarez YD, Fierro-Gonzalez JC, Henon Y, et al. 2015. Cortical tension allocates the first inner cells of the mammalian embryo. *Dev. Cell* 34:435–47
36. Korotkevich E, Niwayama R, Courtois A, Friese S, Berger N, et al. 2017. The apical domain is required and sufficient for the first lineage segregation in the mouse embryo. *Dev. Cell* 40:235–47.e7
37. Totaro A, Panciera T, Piccolo S. 2018. YAP/TAZ upstream signals and downstream responses. *Nat. Cell Biol.* 20:888–99

38. Pocaterra A, Romani P, Dupont S. 2020. YAP/TAZ functions and their regulation at a glance. *J. Cell Sci.* 133:jcs230425
39. Zhao B, Ye X, Yu J, Li L, Li W, et al. 2008. TEAD mediates YAP-dependent gene induction and growth control. *Genes Dev.* 22:1962–71
40. Halder G, Dupont S, Piccolo S. 2012. Transduction of mechanical and cytoskeletal cues by YAP and TAZ. *Nat. Rev. Mol. Cell Biol.* 13:591–600
41. Dupont S, Morsut L, Aragona M, Enzo E, Giullitti S, et al. 2011. Role of YAP/TAZ in mechanotransduction. *Nature* 474:179–83
42. Wada K, Itoga K, Okano T, Yonemura S, Sasaki H. 2011. Hippo pathway regulation by cell morphology and stress fibers. *Development* 138:3907–14
43. Zhao B, Li L, Wang L, Wang CY, Yu J, Guan KL. 2012. Cell detachment activates the Hippo pathway via cytoskeleton reorganization to induce anoikis. *Genes Dev.* 26:54–68
44. Elosegui-Artola A, Andreu I, Beedle AEM, Lezamiz A, Uroz M, et al. 2017. Force triggers YAP nuclear entry by regulating transport across nuclear pores. *Cell* 171:1397–410.e14
45. Panciera T, Azzolin L, Cordenonsi M, Piccolo S. 2017. Mechanobiology of YAP and TAZ in physiology and disease. *Nat. Rev. Mol. Cell Biol.* 18:758–70
46. Ralston A, Rossant J. 2008. Cdx2 acts downstream of cell polarization to cell-autonomously promote trophoblast fate in the early mouse embryo. *Dev. Biol.* 313:614–29
47. Strumpf D, Mao CA, Yamanaka Y, Ralston A, Chawengsaksophak K, et al. 2005. Cdx2 is required for correct cell fate specification and differentiation of trophoblast in the mouse blastocyst. *Development* 132:2093–102
48. Nishioka N, Inoue K, Adachi K, Kiyonari H, Ota M, et al. 2009. The Hippo signaling pathway components Lats and Yap pattern Tead4 activity to distinguish mouse trophoblast from inner cell mass. *Dev. Cell* 16:398–410
49. Niwa H, Toyooka Y, Shimosato D, Strumpf D, Takahashi K, et al. 2005. Interaction between Oct3/4 and Cdx2 determines trophoblast differentiation. *Cell* 123:917–29
50. Frum T, Murphy TM, Ralston A. 2018. HIPPO signaling resolves embryonic cell fate conflicts during establishment of pluripotency in vivo. *eLife* 7:e42298
51. Trimarchi JR, Liu L, Porterfield DM, Smith PJ, Keefe DL. 2000. Oxidative phosphorylation-dependent and -independent oxygen consumption by individual preimplantation mouse embryos. *Biol. Reprod.* 62:1866–74
52. Kaneko KJ, DePamphilis ML. 2013. TEAD4 establishes the energy homeostasis essential for blastocoele formation. *Development* 140:3680–90
53. Kumar RP, Ray S, Home P, Saha B, Bhattacharya B, et al. 2018. Regulation of energy metabolism during early mammalian development: TEAD4 controls mitochondrial transcription. *Development* 145:dev162644
54. Nishioka N, Yamamoto S, Kiyonari H, Sato H, Sawada A, et al. 2008. Tead4 is required for specification of trophoblast in pre-implantation mouse embryos. *Mech. Dev.* 125:270–83
55. Yagi R, Kohn MJ, Karavanova I, Kaneko KJ, Vullhorst D, et al. 2007. Transcription factor TEAD4 specifies the trophoblast lineage at the beginning of mammalian development. *Development* 134:3827–36
56. Chazaud C, Yamanaka Y, Pawson T, Rossant J. 2006. Early lineage segregation between epiblast and primitive endoderm in mouse blastocysts through the Grb2-MAPK pathway. *Dev. Cell* 10:615–24
57. Ohnishi Y, Huber W, Tsumura A, Kang M, Xenopoulos P, et al. 2014. Cell-to-cell expression variability followed by signal reinforcement progressively segregates early mouse lineages. *Nat. Cell Biol.* 16:27–37
58. Schrode N, Saiz N, Di Talia S, Hadjantonakis AK. 2014. GATA6 levels modulate primitive endoderm cell fate choice and timing in the mouse blastocyst. *Dev. Cell* 29:454–67
59. Frankenberg S, Gerbe F, Bessonard S, Belville C, Pouchin P, et al. 2011. Primitive endoderm differentiates via a three-step mechanism involving Nanog and RTK signaling. *Dev. Cell* 21:1005–13
60. Plusa B, Piliszek A, Frankenberg S, Artus J, Hadjantonakis AK. 2008. Distinct sequential cell behaviours direct primitive endoderm formation in the mouse blastocyst. *Development* 135:3081–91
61. Saiz N, Grabarek JB, Sabherwal N, Papalopulu N, Plusa B. 2013. Atypical protein kinase C couples cell sorting with primitive endoderm maturation in the mouse blastocyst. *Development* 140:4311–22

62. Wennekamp S, Mesecke S, Nedelec F, Hiiragi T. 2013. A self-organization framework for symmetry breaking in the mammalian embryo. *Nat. Rev. Mol. Cell Biol.* 14:452–59
63. Rossant J, Tam PP. 2009. Blastocyst lineage formation, early embryonic asymmetries and axis patterning in the mouse. *Development* 136:701–13
64. Ryan AQ, Chan CJ, Graner F, Hiiragi T. 2019. Lumen expansion facilitates epiblast-primitive endoderm fate specification during mouse blastocyst formation. *Dev. Cell* 51:684–97.e4
65. Wang X, Zhang Z, Tao H, Liu J, Hopyan S, Sun Y. 2018. Characterizing inner pressure and stiffness of trophoblast and inner cell mass of blastocysts. *Biophys. J.* 115:2443–50
66. Chan CJ, Costanzo M, Ruiz-Herrero T, Monke G, Petrie RJ, et al. 2019. Hydraulic control of mammalian embryo size and cell fate. *Nature* 571:112–16
67. Weberling A, Zernicka-Goetz M. 2021. Trophectoderm mechanics direct epiblast shape upon embryo implantation. *Cell Rep.* 34:108655
68. Christodoulou N, Weberling A, Strathdee D, Anderson KI, Timpson P, Zernicka-Goetz M. 2019. Morphogenesis of extra-embryonic tissues directs the remodelling of the mouse embryo at implantation. *Nat. Commun.* 10:3557
69. Varmuza S, Prideaux V, Kothary R, Rossant J. 1988. Polytene chromosomes in mouse trophoblast giant cells. *Development* 102:127–34
70. Screen M, Dean W, Cross JC, Hemberger M. 2008. Cathepsin proteases have distinct roles in trophoblast function and vascular remodelling. *Development* 135:3311–20
71. Parast MM, Aeder S, Sutherland AE. 2001. Trophoblast giant-cell differentiation involves changes in cytoskeleton and cell motility. *Dev. Biol.* 230:43–60
72. Bevilacqua EM, Abrahamsohn PA. 1989. Trophoblast invasion during implantation of the mouse embryo. *Arch. Biol. Med. Exp.* 22:107–18
73. Reuss B, Hellmann P, Dahl E, Traub O, Butterweck A, et al. 1996. Connexins and E-cadherin are differentially expressed during trophoblast invasion and placenta differentiation in the rat. *Dev. Dyn.* 205:172–82
74. Zhang T, Guo S, Zhou H, Wu Z, Liu J, et al. 2021. Endometrial extracellular matrix rigidity and IFN γ ensure the establishment of early pregnancy through activation of YAP. *Cell Prolif.* 54:e12976
75. Nieto MA. 2011. The ins and outs of the epithelial to mesenchymal transition in health and disease. *Annu. Rev. Cell Dev. Biol.* 27:347–76
76. Lee K, Nelson CM. 2012. New insights into the regulation of epithelial-mesenchymal transition and tissue fibrosis. *Int. Rev. Cell Mol. Biol.* 294:171–221
77. Savagner P. 2015. Epithelial-mesenchymal transitions: from cell plasticity to concept elasticity. *Curr. Top. Dev. Biol.* 112:273–300
78. Lamouille S, Xu J, Derynck R. 2014. Molecular mechanisms of epithelial-mesenchymal transition. *Nat. Rev. Mol. Cell Biol.* 15:178–96
79. Leight JL, Wozniak MA, Chen S, Lynch ML, Chen CS. 2012. Matrix rigidity regulates a switch between TGF- β 1-induced apoptosis and epithelial-mesenchymal transition. *Mol. Biol. Cell* 23:781–91
80. Lee K, Chen QK, Lui C, Cichon MA, Radisky DC, Nelson CM. 2012. Matrix compliance regulates Rac1b localization, NADPH oxidase assembly, and epithelial-mesenchymal transition. *Mol. Biol. Cell* 23:4097–108
81. Wei SC, Fattet L, Tsai JH, Guo Y, Pai VH, et al. 2015. Matrix stiffness drives epithelial-mesenchymal transition and tumour metastasis through a TWIST1-G3BP2 mechanotransduction pathway. *Nat. Cell Biol.* 17:678–88
82. Brown AC, Fiore VF, Sulchek TA, Barker TH. 2013. Physical and chemical microenvironmental cues orthogonally control the degree and duration of fibrosis-associated epithelial-to-mesenchymal transitions. *J. Pathol.* 229:25–35
83. Matsuzaki S, Darcha C, Pouly JL, Canis M. 2017. Effects of matrix stiffness on epithelial to mesenchymal transition-like processes of endometrial epithelial cells: implications for the pathogenesis of endometriosis. *Sci. Rep.* 7:44616
84. Chen QK, Lee K, Radisky DC, Nelson CM. 2013. Extracellular matrix proteins regulate epithelial-mesenchymal transition in mammary epithelial cells. *Differentiation* 86:126–32

85. Comaills V, Kabeche L, Morris R, Buisson R, Yu M, et al. 2016. Genomic instability is induced by persistent proliferation of cells undergoing epithelial-to-mesenchymal transition. *Cell Rep.* 17:2632–47
86. Simi AK, Anlas AA, Stallings-Mann M, Zhang S, Hsia T, et al. 2018. A soft microenvironment protects from failure of midbody abscission and multinucleation downstream of the EMT-promoting transcription factor Snail. *Cancer Res.* 78:2277–89
87. Rabie EM, Zhang SX, Dunn CE, Nelson CM. 2021. Substratum stiffness signals through integrin-linked kinase and β 1-integrin to regulate midbody proteins and abscission during EMT. *Mol. Biol. Cell* 32:1664–76
88. Copp AJ. 1979. Interaction between inner cell mass and trophectoderm of the mouse blastocyst. II. The fate of the polar trophectoderm. *J. Embryol. Exp. Morphol.* 51:109–20
89. Cockburn K, Rossant J. 2010. Making the blastocyst: lessons from the mouse. *J. Clin. Investig.* 120:995–1003
90. Qin H, Blaschke K, Wei G, Ohi Y, Blouin L, et al. 2012. Transcriptional analysis of pluripotency reveals the Hippo pathway as a barrier to reprogramming. *Hum. Mol. Genet.* 21:2054–67
91. Grapin-Botton A, Constam D. 2007. Evolution of the mechanisms and molecular control of endoderm formation. *Mech. Dev.* 124:253–78
92. Solnica-Krezel L. 2005. Conserved patterns of cell movements during vertebrate gastrulation. *Curr. Biol.* 15:R213–28
93. Kyprianou C, Christodoulou N, Hamilton RS, Nahaboo W, Boomgaard DS, et al. 2020. Basement membrane remodelling regulates mouse embryogenesis. *Nature* 582:253–58
94. Srinivas S, Rodriguez T, Clements M, Smith JC, Beddington RS. 2004. Active cell migration drives the unilateral movements of the anterior visceral endoderm. *Development* 131:1157–64
95. Arnold SJ, Robertson EJ. 2009. Making a commitment: cell lineage allocation and axis patterning in the early mouse embryo. *Nat. Rev. Mol. Cell Biol.* 10:91–103
96. Shioi G, Hoshino H, Abe T, Kiyonari H, Nakao K, et al. 2017. Apical constriction in distal visceral endoderm cells initiates global, collective cell rearrangement in embryonic visceral endoderm to form anterior visceral endoderm. *Dev. Biol.* 429:20–30
97. Migeotte I, Omelchenko T, Hall A, Anderson KV. 2010. Rac1-dependent collective cell migration is required for specification of the anterior-posterior body axis of the mouse. *PLOS Biol.* 8:e1000442
98. Trichas G, Smith AM, White N, Wilkins V, Watanabe T, et al. 2012. Multi-cellular rosettes in the mouse visceral endoderm facilitate the ordered migration of anterior visceral endoderm cells. *PLOS Biol.* 10:e1001256
99. Omelchenko T, Rabadan MA, Hernandez-Martinez R, Grego-Bessa J, KV Anderson, Hall A. 2014. β -Pix directs collective migration of anterior visceral endoderm cells in the early mouse embryo. *Genes Dev.* 28:2764–77
100. Thompson DW. 1917. *On Growth and Form*. Cambridge, UK: Cambridge Univ. Press
101. Park JA, Kim JH, Bi D, Mitchel JA, Qazvini NT, et al. 2015. Unjamming and cell shape in the asthmatic airway epithelium. *Nat. Mater.* 14:1040–48
102. Bi D, Yang X, Marchetti MC, Manning ML. 2016. Motility-driven glass and jamming transitions in biological tissues. *Phys. Rev. X* 6:021011
103. Schotz EM, Lanio M, Talbot JA, Manning ML. 2013. Glassy dynamics in three-dimensional embryonic tissues. *J. R. Soc. Interface* 10:20130726
104. Atia L, Fredberg JJ, Gov NS, Pegoraro AF. 2021. Are cell jamming and unjamming essential in tissue development? *Cells Dev.* In press. <https://doi.org/10.1016/j.cdev.2021.203727>
105. Atia L, Bi D, Sharma Y, Mitchel JA, Gweon B, et al. 2018. Geometric constraints during epithelial jamming. *Nat. Phys.* 14:613–20
106. Wang X, Merkel M, Sutter LB, Erdemci-Tandogan G, Manning ML, Kasza KE. 2020. Anisotropy links cell shapes to tissue flow during convergent extension. *PNAS* 117:13541–51
107. Iyer KV, Piscitello-Gomez R, Paijmans J, Julicher F, Eaton S. 2019. Epithelial viscoelasticity is regulated by mechanosensitive E-cadherin turnover. *Curr. Biol.* 29:578–91.e5
108. Petridou NI, Corominas-Murtra B, Heisenberg CP, Hannezo E. 2021. Rigidity percolation uncovers a structural basis for embryonic tissue phase transitions. *Cell* 184:1914–28.e19

109. Petridou NI, Heisenberg CP. 2019. Tissue rheology in embryonic organization. *EMBO J*. 38:e102497
110. Mongera A, Rowghanian P, Gustafson HJ, Shelton E, Kealhofer DA, et al. 2018. A fluid-to-solid jamming transition underlies vertebrate body axis elongation. *Nature* 561:401–5
111. Lawton AK, Nandi A, Stulberg MJ, Dray N, Sneddon MW, et al. 2013. Regulated tissue fluidity steers zebrafish body elongation. *Development* 140:573–82
112. Kim S, Pochitaloff M, Stooke-Vaughan GA, Campas O. 2021. Embryonic tissues as active foams. *Nat. Phys.* 17:859–66
113. David R, Luu O, Damm EW, Wen JW, Nagel M, Winklbauer R. 2014. Tissue cohesion and the mechanics of cell rearrangement. *Development* 141:3672–82
114. Saadaoui M, Rocancourt D, Roussel J, Corson F, Gros J. 2020. A tensile ring drives tissue flows to shape the gastrulating amniote embryo. *Science* 367:453–58
115. Benazeraf B, Francois P, Baker RE, Denans N, Little CD, Pourquie O. 2010. A random cell motility gradient downstream of FGF controls elongation of an amniote embryo. *Nature* 466:248–52
116. Spurlin JW, Siedlik MJ, Nerger BA, Pang MF, Jayaraman S, et al. 2019. Mesenchymal proteases and tissue fluidity remodel the extracellular matrix during airway epithelial branching in the embryonic avian lung. *Development* 146:dev175257
117. Chambers I, Smith A. 2004. Self-renewal of teratocarcinoma and embryonic stem cells. *Oncogene* 23:7150–60
118. Mitsui K, Tokuzawa Y, Itoh H, Segawa K, Murakami M, et al. 2003. The homeoprotein Nanog is required for maintenance of pluripotency in mouse epiblast and ES cells. *Cell* 113:631–42
119. Nichols J, Zevnik B, Anastasiadis K, Niwa H, Klewe-Nebenius D, et al. 1998. Formation of pluripotent stem cells in the mammalian embryo depends on the POU transcription factor Oct4. *Cell* 95:379–91
120. Takahashi K, Yamanaka S. 2006. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 126:663–76
121. Ramalho-Santos M, Yoon S, Matsuzaki Y, Mulligan RC, Melton DA. 2002. “Stemness”: transcriptional profiling of embryonic and adult stem cells. *Science* 298:597–600
122. Lian I, Kim J, Okazawa H, Zhao J, Zhao B, et al. 2010. The role of YAP transcription coactivator in regulating stem cell self-renewal and differentiation. *Genes Dev.* 24:1106–18
123. Serra D, Mayr U, Boni A, Lukonin I, Rempfler M, et al. 2019. Self-organization and symmetry breaking in intestinal organoid development. *Nature* 569:66–72
124. Nelson CM. 2021. The mechanics of crypt morphogenesis. *Nat. Cell Biol.* 23:678–79
125. Yang Q, Xue SL, Chan CJ, Rempfler M, Vischi D, et al. 2021. Cell fate coordinates mechano-osmotic forces in intestinal crypt formation. *Nat. Cell Biol.* 23:733–44
126. Nelson CM, Gleghorn JP, Pang MF, Jaslove JM, Goodwin K, et al. 2017. Microfluidic chest cavities reveal that transmural pressure controls the rate of lung development. *Development* 144:4328–35
127. Li J, Wang Z, Chu Q, Jiang K, Li J, Tang N. 2018. The strength of mechanical forces determines the differentiation of alveolar epithelial cells. *Dev. Cell* 44:297–312.e5



Contents

Detection and Monitoring of Viral Infections via Wearable Devices and Biometric Data <i>Craig J. Goergen, MacKenzie J. Twardy, Steven R. Steinbubl, Stephan W. Wegerich, Karnika Singh, Rebecca J. Mieloszyk, and Jessilyn Dunn</i>	1
Regulation of Tumor Invasion by the Physical Microenvironment: Lessons from Breast and Brain Cancer <i>Garrett F. Beeghly, Kwasi Y. Amofa, Claudia Fischbach, and Sanjay Kumar</i>	29
Regenerative Approaches for Chronic Wounds <i>François Berthiaume and Henry C. Hsia</i>	61
Current Developments and Challenges of mRNA Vaccines <i>Jinjin Chen, Jianzhu Chen, and Qiaobing Xu</i>	85
Biomaterials for Hemostasis <i>Aryssa Simpson, Anita Shukla, and Ashley C. Brown</i>	111
Direct Cardiac Compression Devices to Augment Heart Biomechanics and Function <i>Jean Bonnemain, Pedro J. del Nido, and Ellen T. Roche</i>	137
Technologies to Assess Drug Response and Heterogeneity in Patient-Derived Cancer Organoids <i>Melissa C. Skala, Dustin A. Deming, and Jeremy D. Kratz</i>	157
Deep Learning and Medical Image Analysis for COVID-19 Diagnosis and Prediction <i>Tianming Liu, Eliot Siegel, and Dinggang Shen</i>	179
Cuffless Blood Pressure Measurement <i>Ramakrishna Mukkamala, George S. Stergiou, and Alberto P. Avolio</i>	203
Derivation and Differentiation of Human Pluripotent Stem Cells in Microfluidic Devices <i>Camilla Luni, Onelia Gagliano, and Nicola Elvassore</i>	231

Improving Antibody Therapeutics by Manipulating the Fc Domain: Immunological and Structural Considerations <i>George Delidakis, Jin Eyun Kim, Katia George, and George Georgiou</i>	249
Cell Trafficking at the Intersection of the Tumor–Immune Compartments <i>Wenxuan Du, Praful Nair, Adrian Johnston, Pei-Hsun Wu, and Denis Wirtz</i>	275
Mechanical Control of Cell Differentiation: Insights from the Early Embryo <i>Celeste M. Nelson</i>	307

Errata

An online log of corrections to *Annual Review of Biomedical Engineering* articles may be found at <http://www.annualreviews.org/errata/bioeng>